

Antibiophilus

Lactobacillus casei

(阿德比膠囊)

組成份：

乳酸桿菌屬(Lactobacillus casei variety rhamnosus)之冷凍乾燥培養物，每公克至少含 8×10^8 個細胞。每-2號膠囊含250公絲。賦形劑含:gelatin,sodium bisulfite,titanium dioxide(E 171)

適應症：

暫時緩解輕度腹瀉、腹痛及便秘、整腸(調整排便)、軟便

注意事項

為避免可能發生的藥物交互作用，應就目前正在使用的任何其他藥物全盤告知醫師或藥師。避免放置於兒童可取用之處。

成人：

治療時併服用甜味飲料及鹹肉湯以補充因腹瀉造成之體液流失。如腹瀉超過2天應請醫師診治。

發生嚴重腹瀉，或糞便內有血或黏液出現，發燒、嘔吐或其他相關的胃腸症狀時應請醫師診治。

兒童：

一嬰兒(2歲以下)：請醫師診治。同時停止餵牛乳及乳製品，給予足量甜味飲料(果汁)及鹹湯(肉湯)以補充因腹瀉造成之體液流失。

每隔6小時測體重。

嬰兒應維持有活力及精神充沛的狀態；否則表示其可能有脫水現象，應立即送醫治療。

—2歲以上兒童：給予本劑外應同時給予再水化治療如甜味飲料及鹹肉湯。

第一天停止給予牛奶及奶製品。

如腹瀉超過兩天應請醫師診治。

發生嚴重腹瀉，糞便內有血或黏液出現，發燒、嘔吐或其他相關的胃腸症狀時應請醫師診治。

其他可能發生的作用

如同其他具療效的藥物一般，本劑亦可能對某些人產生不良作用，必要時請立即向醫師及藥師請教。

用法用量

每天2至8顆。

與液體共服，在2餐之間服用較佳。

請按醫師、藥師、藥劑生指示用藥。

有效期限

請勿超過包裝上所標示之失效日期。

包裝

4~1000粒瓶裝

Laboratoires LYOCENTRE
24, Av. G. Pompidou
15000 AURILLAC-FRANCE



藥商：
旺華生物科技股份有限公司
台北市天母西路13巷6號7樓
TEL:0800-389938

Antibiophilus[®] Capsules

COMPOSITION

Lyophilized culture of Lactobacillus casei variety rhamnosus containing at least 8.10^8 cells per gram.....250mg for one capsule No.2 of 250mg

Composition of capsule tunic:gelatin,sodium bisulfite,titanium dioxide (E 171).

PRESENTATION

Bottle(glass)containing 50 capsules

THERAPEUTIC INDICATIONS

This drug is indicated for the symptomatic treatment of certain diarrhoeas.

The method of rehydration will be determined by your physician.

PRECAUTIONS FOR USE

IN ORDER TO AVOID POSSIBLE DRUG INTERACTIONS, YOU MUST SYSTEMATICALLY INFORM YOUR PHYSICIAN OR YOUR PHARMACIST ABOUT ANY OTHER DRUGS YOU ARE TAKING.KEEP OUT OF THE REACH OF CHILDREN.

In adults:

Add sweetened drinks and salted bouillon to this treatment to compensate for liquid loss due to diarrhoea.

If the diarrhoea lasts longer than 2 days,CONSULT YOUR PHYSICIAN.

In the case of severe diarrhoea, or if blood or mucus is present in the stools,as well as in the case of fever,vomiting or other associated gastrointestinal symptoms,CONSULT YOUR PHYSICIAN.

In children:

- Infants (under 2 years):consult your physician. In the meantime,stop milk and milk-product diet:administer sweetened drinks (fruit juice) and salted drinks(bouillon) in quantities sufficient to compensate for liquid loss caused by diarrhoea. Check bodyweight at 6-hour intervals.

The infant should remain lively and energetic; if this is not the case, it would mean that the infant is dehydrated and should be transferred immediately to hospital

- Children over 2 years:this treatment does not exempt from rehydration:sweetened drinks, salted bouillon, discontinuation of milk and milk-product diet on the first day.

If the diarrhoea lasts longer than 2 day, CONSULT YOUR PHYSICIAN.

In the case of severe diarrhoea, or if blood or mucus is present in the stools, as well as in the case of fever,vomiting or other associated gastrointestinal symptoms, CONSULT YOUR PHYSICIAN.

OTHER POSSIBLE EFFECTS OF THE DRUG

LIKE ANY ACTIVE DRUG, THIS MEDICATION MAY HAVE SOME UNDESIRABLE EFFECTS IN CERTAIN INDIVIDUALS;DO NOT HESITATE TO SEEK ADVICE FROM YOUR PHYSICIAN OR PHARMACIST.

DOSAGE AND METHOD OF ADMINISTRATION

2 to 8 capsules per day.

Swallow the capsule with liquid, preferably not during meals.

IN ALL CASES, FOLLOW YOUR PHYSICIAN'S PRESCRIPTION TO THE LETTER.

CONDITIONS FOR SALE

THIS DRUG WAS PRESCRIBED TO YOU IN A SET OF SPECIFIC CIRCUMSTANCES:

- IT SHOULD NOT BE USED IN ANOTHER SET OF CIRCUMSTANCES
- TREATMENT WITH IT SHOULD NOT BE RESUMED WITHOUT MEDICAL ADVICE
- DO NOT ADVISE SOMEBODY ELSE TO TAKE IT.

SHELF-LIFE

DO NOT EXCEED THE EXPIRY DATE INDICATED ON THE PACKAGE.

Laboratoires LYOCENTRE-24.AV.G.Pompidou-15000 AURILLAC-FRANCE
01077122-03/99 Tél. 04.71.46.80.00